

89011

S/020/60/135/004/004/037  
0111/0222

Some Inequalities for Differentiable Functions of Many Variables

$$(7) I_2 = \left[ \int_{(D_m)} \left( \int_{(D_m)} \left| \frac{\partial^k f(X_m)}{\partial x_{i_1} \dots \partial x_{i_k}} - \frac{\partial^k f(Y_m)}{\partial x_{i_1} \dots \partial x_{i_k}} \right|^q \frac{dY_m}{|X-Y|^{n+\beta q}} \right) dX_m \right]^{1/q} \leq \\ \leq C_9 (A h^{n/q-n/p-k-\beta} + M h^{n-k-\beta}).$$

In (6) and (7),  $h$  is an arbitrary positive number  $\leq 2$ ;  $C_1$  are constants not depending on  $A, M, h, H, \nu, \mu$ .  
For additional assumptions on  $D$  all results can be extended to functions

Card 6/7

07011

8/020/60/135/004/004/037  
0111/C222

Some Inequalities for Differentiable Functions of Many Variables  
having generalized derivatives in the sense of S.L.Sobolev.  
The author mentions S.M.Nikol'skiy and L.N.Slobodetskiy. There are  
4 Soviet references.

ASSOCIATION: Leningradskoye otdeleniye matematicheskogo instituta imeni  
V.A.Steklova Akademii nauk SSSR. (Leningrad Branch of the  
Mathematical Institute imeni V.A.Steklov of the Academy of  
Sciences USSR)

PRESENTED: June 17, 1960, by S.L.Sobolev, Academician

SUBMITTED: June 14, 1960

Card 7/7

16.2600 16.3500

24028  
S/039/61/054/003/001/001  
C111/C444

AUTHOR: Il'in, V. P.

TITLE: Some integral inequalities and their applications in the theory of differentiable functions of several variables.

PERIODICAL: Matematicheskii sbornik, v. 54, no. 3, 1961, 331-380

TEXT: The paper gives a detailed description of the formerly announced results of the author (Ref. 12: Nekotoryye integral'nyye neravenstva dlya differentsirnyemykh funktsiy mnogikh peremennykh [Some integral inequalities for differentiable functions of several variables] DAN SSSR, t. 129, no. 6 (1959), 1214-1217; Ref. 13: Nekotoryye funktsional'nyye neravenstva tipa teorem vlozheniya svesom [Some functional inequalities of the type of embedding theorems with weight] DAN SSSR, t. 129, no. 5 (1959), 983-985). The results overlap partially with V. P. Glushko (Ref. 14: Ob operatorakh tipa potentsiala i nekotorykh teoreмах vlozheniya [On operators of the potential type and some embedding theorems] DAN SSSR, t. 126, no. 3, (1959), 467-470). The author considers generalisations of well-known integral inequalities of the Hilbert inequality type (especially one of a two-parameter inequality of Hardy and Littlewood (Ref. 2: Some properties of fractional integrals. I, Math. Zeitschrift, 27, no. 4 (1928), 565-606)), on Card 1/14

Some integral inequalities...

24028  
S/039/61/054/003/001/001  
C111/444

the higher dimensional case as well as their use in functions of several variables.

Let  $E_n$  be the Euclidean space of  $X = X_n = (x_1, \dots, x_n)$  and  $E_m$  the space of  $Y = Y_m = (y_1, \dots, y_m)$ ,  $1 \leq m \leq n$ . If the coordinates of  $X$  are divided into two groups, then it will be written  $X_n = (X_k, X_{n-k})$ , where  $X_k = (x_1, \dots, x_k)$ ,  $X_{n-k} = (x_{k+1}, \dots, x_n)$ .

Let  $Z = Z_n = (z_1, \dots, z_n)$  be a point of  $E_n$ . We define:  $X_n - Z_n = (x_1 - z_1, \dots, x_n - z_n)$ ;  $Y_m - Z_n = (Y_m - Z_m, Z_{n-m}) = (y_1 - z_1, \dots, y_m - z_m, -z_{m+1}, \dots, -z_n)$ ;

$$r_1 = |X_n - Z_n| = \sqrt{\sum_{i=1}^n (x_i - z_i)^2}, \quad r_2 = |Y_m - Z_n|, \quad r = |X_n - Y_m|.$$

Let  $D = D(X)$  be a measurable set in  $E_n$ ,  $f(X) = f(x_1, \dots, x_n)$  a function, measurable in  $D$  such that  $|f(X)|^p$ ,  $p \geq 1$ , is summable; then is

Card 2/14

24028

S/039/61/054/003/001/001

0111/0444

Some integral inequalities ...

$f \in L_p^{(n)}(D)$  and  $\|f\|_{L_p^{(n)}(D)} = \left[ \int_{D(D)}^{(n\text{-times})} |f|^p dX \right]^{1/p}$ ,  $dX = dx_1 \dots dx_n$ .

$f \in L_{\infty}^{(n)}$  is defined as usual. If  $D = E_n$ , then  $f \in L_p(E_n)$  or  $f \in L_p^{(n)}$  is written instead of  $f \in L_p^{(n)}(E_n)$ . Let  $D(X,Y)$  be a measurable set in  $E_n \times E_m$ .  $D_y(X)$  be the section of  $D(X,Y)$  at a fixed  $Y$ ;  $D(Y)$  being the projection of  $D(X,Y)$  on  $E_m$ , and  $D(X)$  the projection of  $D(X,Y)$  on  $E_n$ . Let  $G(X,Y)$  be a non-negative measurable function in  $D(X,Y)$  and  $f(X)$  measurable in  $D(X)$ .

Let

$$\|fG\|_{L_{p,(n),(m)}} = \sup_{Y \in D(Y)} \left[ \int_{D_y(X)}^{(n\text{-times})} |fG|^p dX \right]^{1/p}$$

if the right hand is finite. In that case be

$fG \in L_{p,\infty}^{(n),(m)}(D(X,Y))$  and if  $D(X,Y) = E_n \times E_m$ , then  $fG \in L_{p,\infty}^{(n),(m)}$ .

Card 3/14

24028  
S/039/61/054/003/001/001  
C111/C444

Some integral inequalities ...

For  $p \geq 1$  let  $p'$  be defined by  $\frac{1}{p} + \frac{1}{p'} = 1$ ; for  $p = \infty$ ,  $q = 1$  and  $q'$  be defined by  $\frac{1}{q} + \frac{1}{q'} = 1$ , let  $\frac{p}{q} = 1$ .

§ 1 Some integral inequalities of the Hilbert inequality type. Five theorems, generalising the above mentioned inequality from [2] in different directions, are proved by aid of four lemmata e. g.

Theorem 1.1:

Let  $p \geq 1$ ,  $q \geq 1$ ,  $\frac{1}{p} + \frac{1}{q} \geq 1$ ,  $1 \leq m \leq n$ ,  $\beta < \frac{m}{q}$ ,  $f(X_n) \in L_p(E_n)$ ,  $g(Y_m) \in L_q(E_m)$ ,  $Z_n = (z_1, \dots, z_n)$ ,  $z_1$  being fixed numbers. Further:

$$1.) K_1 = K_1(X_n - Z_n, Y_m - Z_n) = \begin{cases} r_1^{-\alpha} r_2^{-\beta} r^{\alpha+\beta - (\frac{n}{p} + \frac{m}{q})} & \text{for } r_2 \leq r_1 \\ 0 & \text{for } r_2 > r_1 \end{cases}$$

where  $\alpha > -\beta$  and for  $p > 1$ ,  $\frac{1}{p} + \frac{1}{q} > 1$  holds  $\alpha + \beta \geq 0$

(for  $p = q = 1$ :  $\beta \leq 0$ ,  $\alpha \geq -\beta$ );

$$2.) K'_1 = K_1 r^{\frac{n}{p} + \frac{m}{q} - \lambda}, \text{ where } \lambda < \frac{n}{p} + \frac{m}{q}, \alpha \geq -\beta$$

Card 4/14

24028  
S/039/61/054/003/001/001  
0111/0444

Some integral inequalities ...

(for  $q = 1 : \beta \leq 0$ ),  $h > 0$ .

Then:

$$I_1 = \int_{(E_n \times E_m)}^{n+m} K_1(X_n - Z_n, Y_m - Z_n) f(X_n) g(Y_m) dX_n dY_m \leq M_1 \|f\|_{L_p^{(n)}} \|g\|_{L_q^{(m)}}$$

$$I_1' = \int_{(r \leq h)}^{n+m} K_1'(X_n - Z_n, Y_m - Z_n) f(X_n) g(Y_m) dX_n dY_m \leq$$

$$\leq M_1' \|f\|_{L_p^{(n)}} \|g\|_{L_q^{(m)}} h^{\frac{n}{p} + \frac{m}{q} - \lambda} \quad (17), (17')$$

where  $M_1$  and  $M_1'$  are constants, depending on  $m, n, p, q, \alpha, \beta, \lambda$ , and not depending on  $f, g, h$ .

Theorem 5.1:

1.) Let

$$p > 1, q > 1, \frac{1}{p} + \frac{1}{q} > 1, 1 \leq m < n, 0 \leq \alpha_0 < \frac{n}{p},$$

Card 5/14

Some integral inequalities ...

24028  
S/039/61/054/003/001/001  
C111/C444

$$0 < \beta < \frac{m}{q}, \quad \gamma \frac{p}{q} < \frac{n-m}{p},$$

$$g(Y_m) \in L_q(E_m), \quad f(X_n) |X_{n-m}|^r \in L_p(E_n), \quad f(X_n) r^{-\alpha_0} \in L_{p, \infty}^{(n), (m)},$$

$$K_5 = |Y_m|^{-\beta} r^{\beta + \gamma \frac{p}{q}} - \alpha_0 (1 - \frac{p}{q}) - (\frac{n}{p} + \frac{m}{q}),$$

Then:

$$I_5 = \int_{(E_n, E_m)}^{n+m} K_5 f(X_n) g(Y_m) dX_n dY_m \leq$$

$$\leq K_5 \|g\|_{L_q^{(m)}} \|f |X_{n-m}|^r\|_{L_p^{(n)}} \|f r^{-\alpha_0}\|_{L_{p, \infty}^{(n), (m)}}^{1 - \frac{p}{q}}. \quad (21)$$

Card 6/14



24026  
S/039/61/054/003/001/001  
C111/C444

Some integral inequalities ...

2.) Let  $p > 1$ ,  $q > 1$ ,  $\frac{1}{p} + \frac{1}{q} \geq 1$ ,  $1 \leq m < n$ ,  $0 \leq \alpha_0 < \frac{n}{p}$ ,

$$0 \leq \beta < \frac{m}{q}, (\text{for } q = 1 : \beta = 0), \gamma \leq \frac{n}{q}, (\text{for } p = 1 : \gamma \leq 0)$$

$f$  and  $g$  the same as in 1.),

$$h > 0, \lambda < \frac{n}{p} + \frac{m}{q}, K_5^{(h)} = K_5^{(\frac{n}{p} + \frac{m}{q} - \lambda)}$$

Then:

$$I_5^{(h)} = \int_{(x_n)}^{\dots} \int_{(y_m)}^{\dots} K_5^{(h)} f(x_n) g(y_m) dx_n dy_m \leq \quad (21')$$

$$M_5^{(\frac{n}{p} + \frac{m}{q} - \lambda)} \|g\|_{L_q^{(m)}} \cdot \|f\|_{L_p^{(n)}}^{1-\frac{p}{q}} \|f\|_{L_p^{(n)}, (m)}^{1-\frac{p}{q}}$$

Card 7/14

24028

S/039/61/054/003/001/001  
0111/C444

Some integral inequalities ...

where  $M_5$  and  $M'_5$  depend on  $f$ ,  $g$  and do not depend on  $h$ .

§ 2. Some theorems on integrals of the potential type.

In this paragraph conclusions are brought in eight theorems from the theorems of § 1. e. g.

Theorem 6'. 2: If  $p > 1$ ,  $1 \leq m \leq n$  and besides

$$1) \quad \gamma < \frac{n-m}{p'} ;$$

$$2) \quad \frac{n}{p'} < \lambda + \gamma < \frac{n}{p'} + \frac{m}{p} ;$$

$$3) \quad q' = \frac{m}{\lambda + \gamma - \frac{n}{p'}} ;$$

$$4) \quad |x_{n-m}|^p f(x_n) \in L_p(E_n).$$

then

$$U(Y_m) = \int \overbrace{\dots\dots\dots}^n \frac{f(x_n)}{(x_n - Y_m)} \lambda dx_n \quad L_{q'}(E_m),$$

Card 8/14

Some integral inequalities ...

while  $\|u\|_{L_q(E_n)} \leq M_6 \|f\|_{L_p(E_n)}$ , (29)

holds, where  $M_6$  is independent from  $f$ .

### § 3. Auxiliary inequalities.

#### § 4. Some inequalities for differentiable functions of several variables.

Let  $f(X)$  be a continuous function which is defined in the domain  $D$  of the  $n$ -dimensional space, possessing continuous derivatives in the interior of  $D$ ,  $D$  having the following property: To every point  $P$  of  $D$  one may be able to construct a sector of the  $n$ -dimensional sphere of constant radius and shape, the top of which lies at  $P$ , and which is entirely included in  $D$ . The class of such domains is called  $CD_\alpha$ ,

$\alpha$  being the maximal allowed radius of the sector.

Let 
$$D^j f = \sqrt{\sum_{i_1, \dots, i_j=1}^n \left| \frac{\partial^j f}{\partial x_{i_1} \dots \partial x_{i_j}} \right|^2}.$$

Card 9/14

Some integral inequalities...

Eight theorems are proved e. g.

Theorem 5.4

Let  $1 \leq m < n$ ,  $E_m$  —  $m$  - dimensional hyper-plane in  $E_n$ . Further be

$$p \leq 1, \gamma < \frac{n-m}{p} \text{ (for } p = 1: \gamma \leq 0), 1-k \geq 1,$$

$$1 - k - \gamma > \frac{n-m}{p}$$

$$D \in C_m^n, f(X) \in L_p^{(n)}(D), D^* f \cdot \varphi^v \in L_p^{(n)}(D),$$

where  $\varphi$  is the distance of  $X \in D$  from  $E_m$  and  $q, \delta$  are such that

$$1.) 1 \leq (\gamma, p) \leq q < \infty, \frac{m}{q} > \frac{n}{p} - (1-k) + \gamma,$$

while for  $p > 1$  and  $1-k-\gamma < \frac{n}{p}$  it is  $\frac{m}{q} \geq \frac{n}{p} - (1-k) + \gamma$ .

$$2.) a) \text{ if } 1 - k - \gamma \geq \frac{n}{p}, \text{ then } 0 \leq \beta < \frac{m}{q} \text{ (for } q = \infty: \beta = 0)$$

Card 10/14

21028  
S/039/61/054/003/001/001  
C111/C444

Some integral inequalities ...

b) if  $1 - k - \gamma < \frac{n}{p}$ , then  $0 \leq \beta < \frac{m}{q} - \frac{n}{p} + 1 - k - \gamma$

and for  $p > 1$  it is  $0 \leq \beta < \frac{m}{q} - \frac{n}{p} + 1 - k - \gamma$ .

$D_m$  be the section of  $D$  through  $E_m$ ,  $r_2$  -- the distance of the variable point  $Y \in D_m$  to an arbitrary fixed point of  $D$ .

Then holds the following inequality:

$$\begin{aligned} \|D^k f \cdot r_2^{-\beta}\|_{L_q(D_m)} &\leq K_{10} \|f\|_{L^{(n)}(D)} h^{\frac{m}{q} - \frac{n}{p} + 1 - k - \beta} + \\ &+ K_{11} \|D^1 f \cdot \rho^\gamma\|_{L_p^{(n)}(D)} h^{\frac{m}{q} - \frac{n}{p} + 1 - k - \gamma - \beta}, \end{aligned} \quad (45)$$

where  $h$  is an arbitrary positive number, not exceeding  $k$ , and  $K_{10}$ ,

Carl 11/14

Some integral inequalities ...

$k_{11}$  constants, not depending on  $f$  and  $h$ .

Theorem 6.4

Let  $1 \leq p < \infty$ ,  $0 \leq \alpha_0 < \frac{n}{p}$ ,  $\frac{n}{p} < \alpha < \frac{n}{p}$ ,  $1 - k \leq 1$ .

Let  $D \in C_{\mathbb{R}}^n$ ,  $Z$  a fixed point of  $D$ ,  $f(X) \in L_p(D)$ ,

$v \geq 1$ ,  $r^{\alpha_0} D^1 f \in L_p(D)$ ,  $r^{-\alpha_0} D^1 f \in L_{p, \infty}^{(n), (m)}(D, D)$ ;

the numbers  $m, q, \beta$  satisfy the following conditions:

1)  $1 \leq m \leq n$ ,  $m > -\alpha_p$ ;

2)  $\max(v, p) \leq q \leq \infty$ ;

3)  $\frac{n}{p} - (1 - k) - \frac{\alpha_0}{q} < \frac{m - \alpha_0 p}{q}$ ;

4) a) if  $\frac{n}{p} - (1 - k) - \alpha_0(1 - \frac{p}{q}) + \alpha \frac{p}{q} \leq 0$ , then

Card 12/14

24026  
S/039/61/054/003/001/001  
0111/0444

X

Some integral inequalities ...

21028  
S/039/61/054/003/001/001  
0111/0444

$$-\alpha \frac{p}{q} \leq \beta < \frac{m}{q} \quad (\text{for } q = \infty: \beta = 0)$$

b) if  $\frac{n}{p} - (1 - k) - \alpha_0 (1 - \frac{p}{q}) + \alpha \frac{p}{q} > 0$ , then

$$-\alpha \frac{p}{q} \leq \beta \leq \frac{m}{q} + 1 - k + \alpha_0 (1 - \frac{p}{q}) - \frac{n}{p} - \alpha \frac{p}{q}.$$

Let  $S_m$  be a bounded domain in the intersection  $D_m$  of  $D$  with an  $n$ -dimensional hyperplane. Then the following inequality holds:

$$\| D^k f \cdot r_2^{-\beta} \|_{L_q^{(m)}(S_m)} \leq K_{12} \| D^1 f \cdot r_1^{\alpha} \|_{L_p^{(n)}(D)}^{\frac{p}{q}} \| D^1 f \cdot r^{-\alpha_0} \|_{L_p^{(n), (m)}(D, D_m)}^{1 - \frac{p}{q}} h^{\frac{m}{q} - \frac{n}{p} - k} + K_{13} \| f \|_{L_p^{(n)}(D)}^{\frac{m}{q} - \frac{n}{p} - k} \cdot \begin{cases} h^{-\beta} & \text{for } \beta > 0, \\ d^{-\beta} & \text{for } \beta = 0, \end{cases} \quad (47)$$

Card 13/14

24028  
S/039/61/054/003/001/001  
0111/C444

Some integral inequalities ...

where  $\varepsilon = \frac{m}{n} + \alpha_0 \left(1 - \frac{p}{q}\right) + 1 - k \frac{n}{p} - \frac{p}{q} - \beta,$

$d = \sup_{Y \in S_n} r_2, 0 < n \leq \infty, K_{12} \text{ and } K_{13} \text{ not depending on } f, h, d.$

The author mentions: Kh. L. Smolitskiy.

There are 9 Soviet-bloc and 5 non-Soviet-bloc references. The three references to English-language publications read as follows: G. H. Hardy, J. E. Littlewood, G. Polia, Neravenstva [Inequalities], Moskva, SSL, (1948); G. H. Hardy, J. E. Littlewood, Some properties of fractional integrals. I. Math. Zeitschr., 27, no.4, (1928), 565-606; L. Nirenberg, Estimates and existence of solutions of elliptic equations, Comm. pure and appl. Math., 9, no.3, (1956), 509 - 530.

SUBMITTED: July 4, 1959

Card 14/14



30906

S/517/61/064/000/002/006  
D299/D301

16.4100

AUTHOR: Il'in, V. P.

TITLE: On the approximation of functions of the spaces  $\tilde{W}_p^{(1)}(D)$  and  $W_p^{(1)}(D)$  by continuously differentiable functions

SOURCE: Akademiya nauk SSSR. Matematicheskiy institut. Trudy. v. 64. 1961, 61-78

TEXT: Sufficient conditions are given with respect to the region  $D$  of the  $n$ -dimensional Euclidean space  $E_n$ , under which the functions  $f$ , belonging to Sobolev's spaces  $\tilde{W}$  and  $W$ , can be approximated by continuously differentiable functions. The space  $\tilde{W}_p^{(1)}(D)$  is defined as the totality of functions  $f(X)$  ( $X = x_1, \dots, x_n$ ) on  $D$  which have generalized derivatives (in the sense of S. L. Sobolev) of order  $l = [l]$  ( $[l]$  denoting the integral part of  $l$ ), satisfying the conditions:

Card 1/5

On the approximation ...

S/517/61/064/000/002/006  
D299/D301

$$\|f\|_{L_p(\Omega)} = \left[ \int_{(\Omega)} |f(X)|^p dX \right]^{1/p} < \infty, \quad (1)$$

$$\|f\|_{L_p^{(l)}(\Omega)} = \left[ \sum_{i_1, \dots, i_l=1}^n \left[ \int_{(\Omega)} \left| \frac{\partial^l f(X)}{\partial x_{i_1} \dots \partial x_{i_l}} \right|^p dX \right] \right]^{1/p} < \infty, \quad (2)$$

if  $l$  is an integer, and

Card 2/ 5

S/517/61/064/000/002/006  
D299/D301

On the approximation ...

$$\|f\|_{L_p(D)} = \sum_{i_1, \dots, i_l=1}^n \left\{ \left| \frac{\partial^l f(X)}{\partial x_{i_1} \dots \partial x_{i_l}} \right| \right\} \quad (2')$$

$$\left\{ \left| \frac{\partial^l f(Y)}{\partial x_{i_1} \dots \partial x_{i_l}} \right|^p \cdot \frac{dY}{|X-Y|^{n+(l-1)p}} dX \right\}^{\frac{1}{p}} < \infty,$$

if  $l$  is not an integer; ( $l$  is a non-negative number,  $p \geq 1$ ). The space  $\tilde{W}$  is linear and normalized. A subclass of functions is singled out from the class of functions, belonging to  $\tilde{W}$ ; namely those functions which have, in  $D$ , all possible generalized derivatives to  $l$ -th order inclusive, belonging to  $L_p(D)$ . This subclass is denoted by  $W_p^{(l)}(D)$ . The problem is considered of approximating the functions  $f \in \tilde{W}$  and  $f \in W$ , by a sequence of functions  $\psi_\nu(X)$  ( $\nu = 1, 2, \dots$ ), belonging to  $C^{(1)}(\bar{D})$ , where  $\bar{D} = D + \Gamma$  ( $\Gamma$  being the boundary of  $D$ ).

Card 3/5

On the approximation ...

S/517/61/064/000/002/006  
D299/D301

If  $D = E_n$ , such an approximation can be effected with any degree of accuracy. If, however,  $D$  is not identical with the entire space  $E_n$ , the approximation is not always possible. In the references it was shown that for functions, belonging to  $\tilde{W}$ , the approximation is possible if  $D$  is a bounded region which is starlike with respect to an interior point. In the present work, other sufficient conditions for  $D$  are found, under which the approximation is possible. In particular, the conditions are given, under which the spaces  $\tilde{W}$  and  $W$  coincide. Several lemmas are proved which lead to 3 fundamental theorems. The most important of these, is Theorem 2: Let  $l$  be a non-negative number,  $p \geq 1$ , and  $D$  a region of  $E_n$ , satisfying the conditions: a)  $D$  has the property  $A(N, p)$  (which means that 2 finite systems of  $n$ -dimensional regions  $S_N$  exist, each of which covers  $D$ ); b)  $D$  belongs to  $C(H, \sigma)$  if  $l$  is an integer, and to  $C(H, \sigma, K, \lambda)$ , if  $l$  is not an integer;  $C(H, \sigma)$  means that  $D$  contains an  $n$ -dimensional spherical sector with vertex  $X$ , radius  $H$  and solid

Card 4/5

On the approximation ...

S/517/61/064/000/002/006  
D299/D301

angle  $\sigma$ , for every  $X \in D$ ;  $C(H, \sigma, K, \lambda)$  implies a similar condition with respect to 2 points  $X$  and  $Y$ ). Then: 1) the spaces  $\tilde{W}_p^{(1)}(D)$  and  $W_p^{(1)}(D)$  coincide; 2) for each function  $f \in \tilde{W}$ , there exists a sequence of functions  $\psi_\nu(X)$  which have continuous derivatives of any order in  $E_n$ , so that

$$\lim_{\nu \rightarrow \infty} \|f - \psi_\nu\|_{W_p^{(1)}(D)} = 0.$$

In conclusion, an example is given (taken from the references) which shows that the above approximation is not always possible (as condition  $A(N, \lambda)$  was not satisfied). There are 9 references: 8 Soviet-bloc and 1 non-Soviet-bloc.

Card 5/5

163400

AUTHORS: Il'in, V. P., Solonnikov, V. A.

TITLE: Some Properties of Differentiable Functions of Many Variables

PERIODICAL: Doklady Akademii nauk SSSR, 1961, Vol. 136, No. 3, pp. 538-541

TEXT: The authors consider functions which possess various differential properties in several variables (see (Ref.1-4)).

Let  $f(x_1, \dots, x_n)$  be a smooth function;  $h > 0$ ,  $\alpha_i > 0$  ( $i=1, 2, \dots, n$ ) arbitrary constants;  $v_1, l_1, s_1, k_1$  arbitrary nonnegative integers,  $0 \leq v_1 \leq l_1$ . It holds the identity:

$$D_{x_1}^{v_1} D_{x_2}^{v_2} \dots D_{x_n}^{v_n} f(x) = \frac{C}{K} \int_0^{h_1} \dots \int_0^{h_n} f(x_1 + y_1, \dots, x_n + y_n) \times \\ \times \prod_{j=1}^n \frac{\partial^{l_j}}{\partial y_j^{l_j}} \left[ \frac{y_j^{l_j - v_j - 1}}{(l_j - v_j - 1)!} \psi_j(y_j, h_j) \right] dy_1 \dots dy_n -$$

Card 1/ 10

APPROVED FOR RELEASE

89720  
S/020/61/136/003/003/027  
C 111/ C 333  
Some Properties of Differentiable Functions of Many Variables

$$-C \sum_{i=1}^n x_i \int_{v_i}^{v_i'} \dots \int_{v_1}^{v_1'} \prod_{j=1}^n \frac{\partial f_j}{\partial y_j} \left[ \frac{y_j - v_j}{(v_j - v_j - 1)} \psi_j(y_j, v_j) \right] dy_1 \dots dy_n \times$$

$$\times \int_{v_1}^{v_1'} [D_1 f(x_1 + y_1, \dots, x_1 + y_1 + 1, \dots, x_n + y_n) -$$

$$- 2D_1 f(x_1 + y_1, \dots, x_1 + y_1 + 1, \dots, x_n + y_n) +$$

$$+ D_1 f(x_1 + y_1, \dots, x_1 + y_1 + 1, \dots, x_n + y_n) +$$

$$+ 2y_1 \frac{\partial}{\partial y_1} \int_{v_1}^{v_1'} (v_1' - t)^{r_1 + r_2 + \dots + r_n + v_1} dt,$$

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP

Card 2/ 10

89720  
S/020/61/136/003/003/027  
C 111/ C 333  
Some Properties of Differentiable Functions of Many Variables

$$\psi_j(y_j), v_j' = (1 - v_j) \int_{v_j}^{v_j'} (v_j' - t)^{r_1 + r_2 + \dots + r_n + v_j} dt +$$

$$+ 2y_j \frac{\partial}{\partial y_j} \int_{v_j}^{v_j'} (v_j' - t)^{r_1 + r_2 + \dots + r_n + v_j} dt +$$

$$+ \frac{1}{1 - v_j - 1} y_j \frac{\partial}{\partial y_j} \int_{v_j}^{v_j'} (v_j' - t)^{r_1 + r_2 + \dots + r_n + v_j} dt.$$

Most of the results formulated below follow from the given identity.  
Let D be a domain of the E, with the property: in every point  $y \in \bar{D}$   
an n-dimensional rectangle can be constructed lying entirely in D,  
the corner of which is in x, and the edges of which are parallel to  
the axes of coordinates and have the constant length  $\sum_{j=1}^n (x_j - v_j)$ .  
Moreover: If  $(x_1, \dots, x_1, \dots, x_n)$  and  $(x$

Card 3/ 10

89720

S/020/61/136/003/003/027  
C 111/ C 333

Some Properties of Differentiable Functions of Many Variables

$f \in \mathcal{M}_{1, \dots, 1_n}^{p_0 p_1 \dots p_n}$  can be continued, under remaining differential properties and norm, on the entire  $E_n$  (in the sense of the norm equivalence).

Let

$$\alpha_1 \equiv \frac{1}{l_1} \left( 1 - \sum_{j=1}^n \frac{1}{p_j l_j} + \frac{1}{p_1} \sum_{j=1}^n \frac{1}{l_j} \right) > 0.$$

Theorem 3: Let

$f \in \mathcal{M}_{1, \dots, 1_n}^{p_0 p_1 \dots p_n} (D)$ . Then it holds:

1.) If  $\varepsilon_0 = 1 - \sum_{i=1}^n \frac{1}{p_i l_i} - \sum_{i=1}^n \alpha_i v_i > 0$ , then  $f(x)$  is equivalent to a continuous function differentiable in  $\bar{D}$  and

Card 6/10



89720

S/020/61/136/003/003/027

C 111/ C 333

Some Properties of Differentiable Functions of Many Variables

$$|D_{x_1}^{v_1} D_{x_2}^{v_2} \dots D_{x_n}^{v_n} f| \leq C (\|f\|_{L_{p_0}(D)} h^{-\frac{1}{p_0} \sum_{i=1}^n v_i} + \|f\|_{L_{p_1 \dots p_n}(D)} h^{\epsilon_0}).$$

2.) If  $\epsilon_s = 1 - \sum_{j=1}^n \frac{1}{p_j l_j} - \sum_{j=1}^n \epsilon_j v_j + \frac{1}{q} \sum_{j=1}^s \epsilon_j > 0, q \geq p_1 > 1$   
 then  $(i = 0, 1, \dots, n)$

$$|D_{x_1}^{v_1} D_{x_2}^{v_2} \dots D_{x_n}^{v_n} f|_{L_q(D)} \leq C (\|f\|_{L_{p_0}(D)} h^{\frac{1}{q} \sum_{j=1}^s \epsilon_j - \frac{1}{p_0} \sum_{j=1}^n v_j} + \|f\|_{L_{p_1 \dots p_n}(D)} h^{\epsilon_0}).$$

Card 7/10

89720

S/020/61/136/003/003/027

C 111/ 0 353

Some Properties of Differentiable Functions of Many Variables

3.) If  $q > p_1 > 1$  ( $i = 0, 1, \dots, n$ )

$$1 - \sum_{j=1}^n \frac{1}{p_j^{1/j}} - \sum_{j=1}^n x_j v_j + \frac{1}{q} \sum_{j=1}^n x_j = 0 \text{ then}$$

$$|D_x^\alpha D_x^\beta \dots D_x^\gamma f|_{L_q(D)} \leq C \left( \|f\|_{L_{p_1}(D)} h^{\frac{1}{q} \sum_{j=1}^n \alpha_j - \frac{1}{p_1} \sum_{j=1}^n \beta_j - \frac{1}{p_1} \sum_{j=1}^n \gamma_j} + \|f\|_{L_{p_1 \dots p_n}(D)} \right)$$

$h > 0$  is an arbitrary number which satisfies the condition.

$h \leq H \equiv \min_{i=1, \dots, n} x_i^{1/\alpha_i}$ ,  $D_h$  is the section of  $D$  by the hyperplane

$x_{s+1} = \text{const}, \dots, x_n = \text{const}.$

Card 8/10

89720

S/020/61/136/003/003/027  
C 111/ C 333

Some Properties of Differentiable Functions of Many Variables

Theorem 4 is a further embedding theorem (an analogue of the theorem in (Ref.7) for the Sobolev spaces  $W_p^1$ ).

Theorem 5 says that, if  $D$  is finite, the set  $\{f\}$  is bounded in the norm of

$W_{p_0 p_1 \dots p_n}^{l_1 \dots l_n}(D)$  and the condition 1) or 2) of

theorem 3 is satisfied, then the set

$\left\{ D_{x_1}^{v_1} \dots D_{x_n}^{v_n} f \right\}$  is compact in  $C$  or  $L_q(D_S)$ .

A similar statement refers to theorem 4.

Theorem 6 is a special case of part 3 of theorem 3 and theorem 4, if  $D = E_n$  and  $h = \infty$ .

Card 9/10

89720

S/020/61/136/003/003/027  
O 111/ C 333

Some Properties of Differentiable Functions of Many Variables

Theorem 7 is a partially special inversion of theorem 6.

There are 7 references: 6 Soviet and 1 Italian.

ASSOCIATION: Leningradskoye otdeleniye Matematicheskogo instituta  
imeni V. A. Steklova Akademii nauk SSSR (Leningrad  
Branch of the Mathematical Institute imeni V. A.  
Steklov of the Academy of Sciences USSR)

PRESENTED: July 28, 1960, by V. J. Smirnov, Academician

SUBMITTED: July 21, 1960

Card 10/10

IL'IN V.P.,

Approximation of functions from  $W_p(1)(D)$  and  $W_p(1)(D)$  spaces by continuously differentiable functions. Dokl. AN SSSR 137 no.6:1283-1286 Ap '61. (MIRA 14:4)

1. Leningradskoye otdeleniye Matematicheskogo Instituta imeni V.A. Steklova AN SSSR. Predstavleno akademikom V.I. Smirnovym.  
(Functional analysis)

IL'IN, V.P.; SOLONNIKOV, V.A.

Some properties of differentiable functions of several variables.  
Trudy Mat.inst. 66:205-226 '62. (MIRA 15:11)  
(Functions of several variables)

IL'IN, V.P.

Properties of some classes of differentiable functions of several  
variables defined in an  $n$ -dimensional region. Trudy Mat. inst. 66:  
227-363 '62. (MIRA 15:11)  
(Functions of several variables).

IL'IN, V.P.

Inequalities between the norms of partial derivatives of functions of several variables. Dokl. AN SSSR 150 no. 5:975-977 Je '63.  
(MIRA 16:8)

1. Leningradskoye otdeleniye Matematicheskogo instituta im. V.A. Steklova AN SSSR. Predstavleno akademikom I.M. Vinogradovym.  
(Inequalities (Mathematics)) (Functions of several variables)



IL'IN, V.P.

Some inequalities between the norms of partial derivatives of functions of several variables. Dokl. AN SSSR 152 no.2:262-265.S '63. (MIRA 16:11)

1. Matematicheskii institut im. V.A. Steklova AN SSSR.  
Predstavleno akademikom I.M. Vinogradovym.

IL'IN, V.P.

Splitting of parabolic and elliptic difference equations.  
Sib. mat. zhur. 6 no.6:1425-1428 N-D '65.

(MIRA 18:12)

L 28470-66 ENT(m)/EWA(1)/ENP(r)/TEMP(1)/ETI/EMP(L) TOP(4) JDEM  
ACC NR: AP6010139 (N) SOURCE COOR: UR/0125/61/000/003/0011/0014

AUTHOR: Il'in, V. P.; Razikov, M. I.

ORG: Ural Polytechnic Institute im. S. M. Kirov (Ural'skiy politekhnicheskii institut)

TITLE: Effect of thermal cycle of welding on the properties and phase composition of 30Kh10G10 chromium-manganese steel

SOURCE: Avtomaticheskaya svarka, no. 3, 1966, 11-14

TOPIC TAGS: austenitic steel, chromium steel, manganese steel, arc welding, thermal effect, cooling rate, phase composition / 30Kh10G10 austenitic steel

ABSTRACT: 30Kh10G10 austenitic metastable chromium-manganese steel is highly cavitation-resistant and finds application as a protective coating for low-carbon and low-alloy metal of the parts of hydraulic machinery and water turbines. When heated above 600°C, however, its structure becomes unstable, which affects its mechanical properties. Hence it was of interest to investigate the thermal effect of welding cycle on the phase composition and mechanical properties of the metal in the zone affected by the welding heat. Specimens of this steel were tested under conditions simulating the effect of the thermal cycle of single-pass arc welding; the maximum

Card 1/2

UDC: 621.791.01:669.140.56.046

ACC NR: AP6010139

temperature was 1350-1400°C and the cooling rate varied within the limits of from 600 to 0.15°C per sec in the 750-650°C range. In this case the time of stay of specimens of this steel (3-5 mm thick sheets) in the dangerous temperature range (800-600°C) is determined by their cooling rate. The lower the cooling rate is, the greater the effect of the thermal cycle on the mechanical properties and phase composition of the metal. Findings: at a cooling rate of less than 1°C/sec the impact strength of the specimens drops sharply, although their ultimate strength remains almost the same whether the cooling rate is 600°C/sec or 0.15°C/sec. At cooling rates of from 600 to 40°C/sec hardness changes insignificantly and amounts to ~250 kg/mm<sup>2</sup>, but if the cooling rate is further reduced (to 1.2°C/sec), hardness begins to increase and reaches 290 kg/mm<sup>2</sup>, but below 1.2°C/sec it decreases again, evidently owing to the conglomeration and precipitation of fine-dispersed carbides. Radiographic examination showed that the decrease in the impact strength of 30Kh10G10 steel is due to both the decomposition of austenite with segregation of  $\gamma$ - and  $\alpha$ -phases and the precipitation of carbides at grain boundaries with consequent embrittlement of grain boundaries. These structural transformations ( $\gamma \rightarrow \epsilon \rightarrow \alpha$ , with ultimate segregation of carbide phase) account for the deterioration in the plasticity and impact strength of the metal. Thus, the welding regime of 30Kh10G10 steel should be such as to assure a cooling rate of at least 1°C/sec; hence, during single-pulse welding of 30Kh10G10 sheet steel (3-5 mm thick) the linear energy should not exceed 2000-2500 cal/cm, i.e. this steel should be welded at minimal current intensity and high speed. Orig. art. has: 4 figures, 3 tables.

SUB CODE: 11, 13/ SUBM DATE: 30Nov64/ ORIG REF: 004

Card 2/2 LC

IL'IN, V.P.

Some properties of classes of differentiable functions defined  
in a certain region. Trudy Mat. inst. 84:93-113 '65.

Inequalities between the norms of partial derivatives of functions  
of several variables. Ibid.:144-173 (MIRA 18:9)

IL'IN V. S.

"The Structure and Functional-Mechanical Properties of the Wood of Various Forms of Birch as Distinguished by Its Bark." Cand Agr Sci, Leningrad Order of Lenin Forestry Engineering Acad imeni S. M. Korov, Leningrad, 1954. (KL, No 7, Feb 55)

SO: Sum. No. 631, 26 Aug 55- Survey of Scientific and Technical Dissertations Defended at USSR Higher Educational Institutions (14)

IL'IN, V. S.

USSR/Radio Receivers, Battery  
Radio receivers - Measurements

Jan 1946

"Utilization of Receiving Apparatus of Radio Centers  
for Two-way Group Telephone Communication," V. S.  
Il'in, Engineer at the Telegraph-Telephone Section  
of the Ministry Administration of the People's  
Commissariat of Communications, 1 p

"Vestnik Svyazi - Elektro Svyaz'" No 1 (70)

Use of CTe-6-M apparatus showed a lack of volume  
over long distance communication. Amplifier  
changes would entail too much new apparatus and  
an uneconomical expenditure of fuel. I. S. Smolin  
developed a method by which the standard battery  
19742

USSR/Radio Receivers, Battery (contd) Jan 1946  
Radio receivers - Measurements

receivers RI-234 and RPK-9 and others could be so  
utilized to carry out two-way amplification of  
the flow of low frequencies.

19742

IL'IN, V.S.

Using the variation method for calculations of "sudden" inhomogeneities in wave guides. Izv.vys.ucheb.zav.; radiofiz. 1 no. 188-94 '58.  
(MIRA 12:1)

1. Saratovskiy gosudarstvennyy universitet.  
(Wave guides)



85988

S/141/60/003/004/013/019  
E192/E382


9,1300 (also 1006)

AUTHOR: Il'in, V.S.

TITLE: Variation Method of Calculating the Symmetrical  
Quadripole Discontinuities in Rectangular Radio  
Waveguides 25PERIODICAL: Izvestiya vysshikh uchebnykh zavedeniy,  
Radiofizika, 1960, Vol. 3, No. 4, pp. 683 - 693

TEXT: The Schwinger variation method was employed by Il'in (Ref. 1) to analyse the discontinuity encountered when joining two rectangular waveguides of different heights. In the following the method is extended to the case when the waveguides are coupled by means of a grid-like diaphragm having a finite length. The system considered is shown in Fig. 1. The grid-like discontinuity has a thickness  $L$ . It is assumed that in the axis  $y$  (perpendicular to the plane of the drawing), the system is uniform and is bounded by two walls situated in the planes  $y = 0$  and  $y = d$ . Consequently, the problem should be solved for the E-plane of the waveguide. Moreover, it is assumed that the losses in the media filling the waveguides I and II, the channels of the discontinuity


Card 1/5



85988

S/141/60/003/004/013/019  
E192/E382Variation Method of Calculating the Symmetrical Quadripole  
Discontinuities in Rectangular Radio Waveguides

and the walls are negligible. The spacings  $b$  (Fig. 1) are chosen so that only the dominant waves of the type  $H_{0m}$  are propagating in the two waveguides. The electric field components  $E_x$  in the regions 1, 2, 3, ..., I and II can therefore be expressed by Eqs. (1), where the propagation constants are defined by Eqs. (2).  $R$  and  $T$  in the above equations represent the reflection and transmission coefficients with respect to the terminal planes  $z = 0$  and  $z = L$ . By comparing the field components at  $z = 0$ , a set of formulae represented by Eqs. (3) is obtained. A similar procedure for  $z = L$  leads to Eqs. (4). The magnetic field component  $H_y$  can be determined from Eq. (5). The comparison of the magnetic field components at  $z = 0$  and  $z = L$  leads to a system of two integral equations which are expressed by Eqs. (6) and (7).



Card 2/5

85988

S/141/60/003/004/013/019

E192/E382

Variation Method of Calculating the Symmetrical Quadripole Discontinuities in Rectangular Radio Waveguides

These can be transformed by addition and subtraction so that two different integral equations are derived; these are given by Eqs. (8). Further transformation of Eqs. (8) can be effected by introducing the notation defined by Eqs. (9). From Eqs. (3), (4) and (9), it is possible to derive Eq. (10).

By multiplying Eqs. (8) by  $E^{-(+)}$  and then integrating them over 0 to b and taking into account Eq. (10), the relationship between R and T can be expressed by Eqs. (11). These represent two quadratic functionals. The problem now consists of finding stationary points of the functionals given by Eqs. (11). In other words, it is necessary to solve a variation problem. The functions  $E^{-(+)}$  can be represented in the form of the series given by Eqs. (12), where the coefficients a can be defined from the conditions expressed by Eqs. (13). The functions  $E^{-(+)}$  can be approximately expressed by Eq. (14). Consequently, the functionals can be

Card 3/5

8 5988

S/141/60/003/004/013/019

E192/E382

Variation Method of Calculating the Symmetrical Quadripole Discontinuities in Rectangular Radio Waveguides

written in the form of Eqs. (15). It is now possible to determine the equivalent circuit of the discontinuity. This is represented by the T-type network shown in Fig. 2. The impedances of this network are expressed by:

$$Z = \frac{1 + R - T}{1 - R + T} ; Z + 2Z_{\text{ш}} = \frac{1 + R + T}{1 - R - T} \quad (16) .$$

On the other hand, by comparing Eqs. (15) and (16) it is found that the impedances can be determined from Eqs. (17). Eqs. (11) and (15) are employed to analyse some special cases. First, the case of  $L = 0$  is investigated (Fig. 3a). Other cases which can be analysed by employing the above method are illustrated by the remaining sketches in Fig. 3. It is finally pointed out that a similar method can be employed to analyse the H-plane in waveguide systems.

Card 4/5

85988

S/141/60/003/004/013/019

E192/E382

Variation Method of Calculating the Symmetrical Quadripole Discontinuities in Rectangular Radio Waveguides

This paper was presented at the Third All Union Conference MVO SSR on Radioelectronics.

There are 3 figures and 7 Soviet references, 1 of which is translated from English.

ASSOCIATION: Saratovskiy gosudarstvennyy universitet  
(Saratov State University)

SUBMITTED: March 19, 1960

Card 5/5

KALININ, V.I., prof., doktor fiziko-matem. nauk [deceased];  
AKINDINOV, V.V.; GERSHTEYN, G.M.; DASHENKOV, V.M.; YEVSEYEV,  
V.I.; IL'IN, V.S.; KOROSTELEV, G.N.; LUCHININ, V.D.; MAJMEIKO,  
Yu.P.; RYAZANOVA, T.P.; SEDIN, V.A.; TOLSTIKOV, V.A.; SHITYROV,  
A.I.; AVILOV, B.I., red.; ZENIN, V.V., tekhn. red.

[Practical work in radio physics] Radiofizicheskiy praktikum.  
Izd.2., dop. i perer. Saratov, 1961. 277 p. (MIRA 15:1)

1. Saratov. Universitet. 2. Kafedra radiofiziki Saratovskogo  
universiteta im. N.G.Chernyshevskogo (for all except Avilov,  
Zenin).

(Radio)

IL'IN, V.S.; PITOVA, G.V.

Sulfhydryl groups of yeast hexokinase: their role in catalysis and in hormone action on enzyme activity. Vop.med.khim. 10 no.2:214-216 Mr-Apr '64. (MIRA 18:1)

2. Otdel biokhimi Instituta eksperimental'noy meditsiny AMN SSSR, Leningrad.

is known. The method is set forth as applied to four-pole irregularities in

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618510005-5

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618510005-5"

5/0109/65/010/001-0-01/281

AUTHOR: Darbenkov, V. M. & Il'In, V. G.

... axially symmetrical delay structures

... 1965, 1-2, 293

... axially symmetrical delay structures

... method ... applied to a wide ...  
... axially symmetrical structure with ...  
... of electric field  $E_z$ ,  $E_y$  ...  
... by the Ritz method ...  
... The method is applied ...  
... delay structures ...



... systems. (c) a second system, (d) a ...  
... selection. TW-tube in BW series the ...  
... method. The dispersion equation is ...  
... with respect to the propagation ...  
... The cumbersome calculation ...  
... Orig. has a dispersion ...

APPROPRIATE/AM-1000

... ..

... ..

... ..

SUBJECT: ...

... ..

... ..

KULIKOV, E.L.; IL'IN, V.S.; IL'IN, V.K.

Design of irregular waveguides using the "equivalent resonator"  
method. Izv. vys. ucheb. zav.; radiotekh. 8 no.1:18-26 Ja-F '65.  
(MIRA 18:5)

DASHENKOV, V.M.; IL'IN, V.S.

Variational method for calculating axisymmetrical collay systems.  
Radiotekh. i elektron. 10 no.2:269-283 F '65.

(UKHA 18:3)

IL'IN, V.S.; STEPANOVA, N.G.

Activity of hexokinase, glucose-6-phosphate dehydrogenase and glucose-6-phosphatase of liver cell fractions in rabbit embryos and full-grown animals following introduction of glucocorticosteroids. Vop. med. khim. 10 no.6:576-584 N-D '64.

(MIRA 19:1)

1. Otdel biokhimii Instituta eksperimental'noy meditsiny AMN SSSR, Leningrad.

DOKUSOVA, O.K.; IL'IN, V.S.

Changes in energy metabolism in the liver of chickens with  
biotin deficiency. Biokhimiia 29 no.5:854-862 J1-Ag '64.  
(MIRA 18:11)

1. Otdel biokhimii Instituta eksperimental'noy meditsiny AN  
SSSR, Leningrad.

IL'IN, V.S.; TITOVA, G.V.

Hormone-enzyme complex: insulin-hexokinase. Biokhimiia  
30 no.6:1251-1256 N-D '65. (MIRA 19:1)

1. Otdel biokhimi Institute eksperimental'noy meditsiny  
AMN SSSR, Leningrad. Submitted April 3, 1965.

L 2608-66 EWT(1)/ZWA(h)

ACCESSION NR: AP5020134

UR/0109/65/010/008/1540/1542

621.517.34

AUTHOR: Dashenkov, V. K.

TITLE: Measuring the coupling resistance of delay systems by means of dielectric bars

SOURCE: Radiotekhnika i elektronika, v. 10, no. 8, 1965, 1540-1542

TOPIC TAGS: delay system 25

ABSTRACT: A measuring technique is suggested which permits easy determining of squared spatial harmonics of the longitudinal and transverse components of the standing-wave electric field  $E_z^2$  and  $E_x^2$  for substituting them into this formula  $\xi = \alpha_z E_z^2 + \alpha_x(\theta) E_x^2$ . The formula gives the relative detuning of a delay system caused by insertion of a thin isotropic dielectric bar;  $\alpha_z$  and  $\alpha_x$  are the bar polarizabilities;  $\theta$  is the angle between the transverse component  $E_x$  and the x-axis. In order to determine the above spatial harmonics, it is recommended to measure the maximum and minimum detunings for perpendicular orientations of the probe with respect to  $E_x$ . The formulas were verified experimentally on an interdigital delay system consisting of 32 studs into which a

Card 1/2

L 2608-66

ACCESSION NR: AP5020134

0.51 x 3.51-mm ebony strip was introduced. Orig. art. has: 2 figures, 12 formulas, and 1 table.

ASSOCIATION: none

SUBMITTED: 18Nov64

ENCL: 00

SUB CODE: 120

NO REF SOV: 004

OTHER: 000

Card

2/2



L 27677-66 ENP(k)/ENT(1)/T GG

ACC NR: AP600752

SOURCE CODE: UR/0141/66/009/001/0016/0032

AUTHOR: Il'in, V. S.; Kostyunina, O. P.

ORG: Saratov State University (Saratovskiy gosudarstvennyy universitet)

TITLE: Diffraction of electromagnetic waves by ultrasonic waves in anisotropic medium -- Part 1

SOURCE: IVUZ. Radiofizika, v. 9, no. 1, 1966, 16-32

TOPIC TAGS: electromagnetic wave diffraction, ultrasonic diffraction

ABSTRACT: Diffraction patterns have been calculated by solving a scalar problem for the Fraunhofer zone; space effects in the medium, interface-reflection effects, etc. have not been fully accounted for (P. Debye, Ber. Sach. Akad. Wiss., v. 84, 125, 1932; C. Raman, N. Nath, Proc. Ind. Inst. Sc., 3(A), 75, 1936). The present article uses the variational method for studying the transmission of electromagnetic waves through an anisotropic layer (a crystal) whose dielectric constant is modulated by ultrasonic waves. Formulas for the diffraction field in both near and remote zones are developed. It is proven that, for the near zone, the system can be described by a diffusion matrix  $S$  of a  $2(N+1)$ -terminal network; this permits using the matrix method of circuit theory for studying various cascade connections of such devices. The above formulas cover, as particular cases, the classical formulas of scalar diffraction in an isotropic layer. Orig. art. has: 1 figure and 90 formulas.

SUB CODE: 20, 09 / SUBM DATE: 19Jul65 / ORIG REF: 009 / OTH REF: 007

Card 1/1

UDC: 621.371.167

L 31916-66 ENT(1)/T WR

ACC NR: AF6010726

SOURCE CODE: UR/0142/66/009/001/0081/0089

AUTHOR: Il'in, V. S.


ORG: none

TITLE: Reflection properties of compound metal-dielectric arrays

SOURCE: IVUZ. Radiotekhnika, v. 9, no. 1, 1966, 81-89

TOPIC TAGS: ~~THE~~ ANTENNA; ANTENNA ARRAY; DIELECTRIC PROPERTY; ELECTROMAGNETIC WAVE REFLECTION

ABSTRACT: General approximate formulas are developed for the reflection and transmission coefficients of a planar wave normally falling on a laminated dielectric slab reinforced with shaped metal parts (see figure). The formulas are in the integral form and are



UDC: 621.396.673.4

Card 1/2

L 27660-66

ACC NR: AP6008184

SOURCE CODE: UR/0109/00/011/003/0411/0476

AUTHOR: Dashenkov, V. M.; Il'in, V. S.; Navrotskaya, Yu. M.

ORG: none

TITLE: Calculating natural frequencies of axially-symmetrical resonators and critical wavelengths of regular waveguides

SOURCE: Radiotekhnika i elektronika, v. 11, no. 3, 1966, 471-476

TOPIC TAGS: resonator, waveguide

ABSTRACT: The field structure in some periodic waveguides is such that description of the delay system, with  $\varphi = 0$  and  $\varphi = \pi$ , can be reduced to consideration of the resonators whose configuration is determined by the geometry of one section of the delay system. Hence, the delay-system dispersion equation can be used for calculating the natural frequencies of such resonators. In an extreme case, when the radii of the delay system approximate infinity, the axially-symmetrical resonator "degenerates" into a shaped regular waveguide. Authors' formulas developed earlier (Rad. i elektronika, 1965, v. 10, no. 2, 269) for axially-symmetrical delay systems are adopted for single-ridge waveguides; this method is illustrated by a calculation of fundamental TE-mode in such a waveguide. Orig. art. has: 3 figures, 35 formulas, and 1 table.

SUB CODE: 09 / SUBM DATE: 18Nov64 / ORIG REF: 004 / OTH REF: 000

UDC: 621.372.834.001.24


Card 1/1

CR

117

The glutathione content of the tissues and the blood after transfusion of large amounts of blood from another species. V. S. Illin. *Bull. Biol. Med. Acad. U. S. S. R.* 6, 648-650 (1968). *Chem. Zvest.* 1968, 11, 2251-4. After the administration of large amounts of blood of a foreign species the glutathione content was increased in the kidneys, muscles and especially in the liver. This increase was chiefly at the expense of the sulphydryl form. No essential change in the glutathione content of the heart muscle was observed. Extensive changes in the glutathione content of the blood took place, although no strict regularity in such changes in the blood could be established. According to Illin, the changes in glutathione content occurring in the tissues are due to disturbances of other processes of tissue metabolism and above all to those in protein metabolism. From this standpoint it is of interest to note that the greatest glutathione increase takes place directly in the liver, which is known to be the seat of the most intensive proteolytic processes in the organism.

M. G. Mikov



1ST AND 2ND COLUMNS		PROCESSING AND REPRODUCTION INDEX		3RD AND 4TH COLUMNS	
ca.				119	
<p>The oxidation coefficient of the urine and blood of dogs after injection of heterogenous blood. V. S. Il'in. <i>Bull. Acad. Sci. USSR Div. Med. Sci. Biol. Sci.</i> 1958, 4: 475-4 (1958) (in English).—Injection of heterogenous blood increased the <math>\text{variet-O}</math> of the blood. If the increase of sugar and lactic acid is taken into consideration, it becomes evident that the change of <math>\text{variet-O}</math> is insignificant and, consequently, the oxidation coeff. of the blood is practically unchanged. On injection of 3-5 cc./kg. of human blood into dogs no noticeable increase of the immediately oxidized products in the blood was observed, and no increase of excretion of such products by the kidneys was obtained. Dogs without morphine endure the injection of heterogenous blood better than those under morphine narcosis. Morphine narcosis seems to be one of the factors increasing the sensitivity of the dogs to heterogenous blood. W. H. H.</p>					
A 58-543 METALLURGICAL LITERATURE CLASSIFICATION		U.S. DEPT. OF COMMERCE		U.S. DEPT. OF COMMERCE	
15000 51700000		15000 51700000		15000 51700000	
15000 51700000		15000 51700000		15000 51700000	

<p>ca.</p> <p>Effect of hormones of the adrenal cortex and medulla on the glutathione content of blood. V. S. H'm. <i>J. Physiol.</i> (U.S.S.R. 34, 862 U in English, 1961, 1961), of 1961 34, 2439. - The glutathione (G) content of liver, kidney and blood of rats is only slightly increased by adrenocortical, as by the removal of one and demedullation of the other adrenal gland. Adrenaline, noradrenaline, and cortisol hormone decreases the G of the muscles. T. Langer.</p>	
<p>ASB. 5.5.5 METALLURGICAL LITERATURE CLASSIFICATION</p>	
<p>ADONIS 0000-0000</p>	<p>ADONIS 0000-0000</p>
<p>ADONIS 0000-0000</p>	<p>ADONIS 0000-0000</p>





IL'IN, V. S.

FA 1/24/74

USSR/Medicine - Fibrinogenase  
Fibrin, Fibrinogen

Nov/Dec 1953

"Fibrinogenase," V. S. Il'in, Leningrad, 94 pp.

"Uspekhi Sovrem Biol" Vol XXVI, No 3 (6)

Tests with blood obtained from corpses and from living organisms proved that fibrinogenase does not convert fibrinogen to fibrin; Transition occurs in another albumin (or other albumins) and there is no coagulation under the influence of thrombin. Name "Fibrinogenodestructase" is considered more proper on strength of above tests.

3/5044

1ST AND 2ND SERIES

PROCESSES AND PROPERTIES

ca

11f

The fibrinogen and fibrinolytic factors (fibrinogenase) of blood of people who meet sudden death. V. B. M. Stachursky (4, 514-80(1940); cf. Wobbsch and Jähling, C. A. B. 5223. The blood of a person who died suddenly not only does not coagulate, but it possesses the property of liquefying coagulated blood of normal people and animals. The fibrinogenase and fibrinolytic factor of such blood is thermostable, and is completely destroyed on heating at 60-65° for 30 min. An extract of fibrinogenase can be prepared from the lungs of people who died suddenly. The active fibrinogenase leaves the lungs and enters the body during sudden death. H. Stachursky

ALL INFORMATION CONTAINED HEREIN IS UNCLASSIFIED

DATE 11-11-80 BY 1045

1045

ILIN, V. S.

U.S.S.R. / Human Animal Physiology. Metabolism. T

Abs Jour: Ref Zhur-Biol., No 5, 1958, 21837.

Author : ~~Ilia V. S.~~


Inst : Institute of Experimental Medicine-Academy of  
Medical Sciences, U.S.S.R.

Title : Regulation of Energy Metabolism and Its Mechanism.

Orig Pub: Yezhegodnik-Inst. Experim. Med. Nauk 1955,  
L 1956, 189-204.

Abstract: Resume: The significance of the process of oxidizing phosphorylation and of different enzyme systems partaking in this process was considered in reference to energy metabolism of the cell. Data were furnished concerning changes of these factors by elevation of body temperature and starvation.

Card 1/1



*I L 50, 4.0.*  
VESELKIN, N.V.; DAUDOVA, G.M.; IL'IN, V.S.

Method of determining the ability of tissues to synthesize reducing  
phosphorus organic compounds. Met. po evol. fiziol. 1:76-78 '56.  
(PHOSPHORUS ORGANIC COMPOUNDS) (MIRA 11:1)  
(TISSUE CULTURE)

USSR / Human and Animal Physiology (Normal and Pathological). Blood. Blood Pressure. Hypertonia T

Abs Jour: Ref Zhur-Biologiya, No 21, 1958, 97574

Author : Veselkin, N. V., Il'in, V. S.

Inst : Not given

Title : Synthesis of Ethero-Sulfuric Acids in an Organism After Denervation of the Liver

Orig Pub: V Sb.: Materialy po evolyuts. fiziologii, T. I. M.-L., AN SSSR, 1956, 85-90

Abstract: No abstract

Card 1/1

41

IL'IN, V.S.; TITOVA, G.V.

Mechanism of the action of insulin on the hexokinase reaction.  
Vop.med.khim. 2 no.3:203-209 My-Je '56. (MLRA 9:10)

1. Otdel biokhimii Instituta eksperimental'noy meditsiny ANS SSSR,  
Leningrad

(INSULIN, effects,  
on hexokinase (Rus))

(TRANSPHOSPHORYLASES,  
hexokinase, eff. of insulin (Rus))

IL'IN, Y.S.; TITOVA, G.V.

Relation of hexokinase-inhibiting properties of  $\beta$ -lipoprotein fraction of blood serum to cortisone and insulin. Vop.med.biol. 2, no. 4: 243-251 J1-Ag '56. (MIRA 9:10)

1. Otdel biokhimi i insituta eksperimental'noy meditsiny AN SSSR, Leningrad.

(BLOOD PROTEINS,

$\beta$ -lipoprotein fractions, inhib. of hexokinase, eff. of cortisone & insulin (Rus))

(TRANSPHOSPHORYLASES,

hexokinase, inhib. by blood  $\beta$ -lipoprotein fractions, eff. of cortisone & insulin (Rus))

(CORTISONE, effects,

blood  $\beta$ -lipoprotein fractions inhib. of hexokinase (Rus))

(INSULIN, effects,

same)



V.S. ILIN  
USSR/RS  
Abstr Jour: *Izv. Akad. Nauk SSSR Ser. Biol.*  
Author: *Y.A. Il'in, T.I. Vol'fson, Z.A. Chuplygin and K.F. Kravchenko*  
Inst Title: *The Influence of the Nervous System on the Activity of Blood Fibrinogenase.*  
Orig Pub: *Tr. Vses. obschestva fiziol., biohim. i farmakologov, 1956, 3, 117-118.*

Abstract: Active fibrinogenase was not detected in the blood of 30 healthy individuals, but was found in the blood of 22 out of 40 surgical patients on the day preceding a serious operation. In these same patients the active enzyme was found in only five cases a day after the operation. Analogous data was obtained

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618510005-5

Card : 1/3

Card : 2/3

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618510005-5

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618510005-5"

VESELIKIN, N.V.; DAUDOVA, G.M.; IL'IN, V.S.

Acid-soluble phosphorus fractions in the mammary glands of rabbits  
and the activity of glands in synthesizing reducing phosphorus  
organic compounds. Mat. po evol. fiziol. 1:79-84 '57. (MIRA 11:1)  
(MAMMARY GLANDS)  
(PHOSPHORUS ORGANIC COMPOUNDS)

IL'IN, V.S.

Significance of hexokinase reaction and mechanisms regulating  
it. Ukr.biokhim.zhur. 30 no.6:911-943 '58. (MIRA 11:12)  
(HEXOKINASE)

IL'IN, Vasil'y S., otv.red.; BIRYUKOV, D.A., red.; MEYFAKH, S.A., red.

[Phosphorylation and its functions; the 1958 symposium] Fosfori-  
lirovanie i funktsiia; simposium 1958 g. Leningrad, 1960. 377 p.  
(MIRA 14:7)

1. Akademiya meditsinskikh nauk SSSR, Moscow. Institut eksperi-  
mental'noy meditsiny.  
(Phosphorylation)

KHARAUZOV, N.A., prof., glavnyy red.; MIKHAYLOV, V.P., prof., zamestitel' glavnogo red.; BIRYUKOV, D.A., prof., otv.red.; AVER'KIN, B.G., doktor biol.nauk, red.; ANICHKOV, N.M., akademik, red.; ANICHKOV, S.V., prof., red.; ARBUZOV, S.Ia., prof., red.; VESSELKIN, P.M., prof., red.; VOYNO-YASENETSKIY, M.V., prof., red.; DANILOV, I.V., kand.biol.nauk, red.; ZHABOTINSKIY, Yu.M., prof., red.; ZHINKIN, L.N., prof., red.; IL'IN, V.S., red.; IOFFE, V.I., prof., red.; KARASIK, V.M., prof., red.; KUPALOV, P.S., prof., red.; MANINA, A.A., kand.med.nauk, red.; NEMYFAKH, S.A., doktor biol.nauk, red.; RIKKL', A.V., prof., red.; SVETLOV, P.G., prof., red.; SMORODINTSEV, A.A., prof., red.; CHISTOVICH, G.N., doktor med.nauk, red.; BESKUDIN, I.K., tekhn. red.

[Yearbook of the Institute of Experimental Medicine of the Academy of Medical Sciences of the U.S.S.R. for 1958] Ezhegodnik za 1958 god. Leningrad, 1959. 538 p. (MIRA 14:1)

1. Akademiya meditsinskikh nauk SSSR, Moscow, Institut eksperimental'noy meditsiny. 2. Chleny-korrespondenty Akademii meditsinskikh nauk SSSR (for Biryukov, Vesselkin, Il'in, Ioffe, Karasik, Svetlov, Smorodintsev). 3. Deyatvitel'nyye chleny Akademii meditsinskikh nauk SSSR (for Anichkov, S.V., Kupalov).

(MEDICINE, EXPERIMENTAL)

IL'IN, V.S.; MEYFAKH, S.A.

Conference on the problem of "Phosphorylation and function."  
Vop.med.khim. 5 no.2:153-159 Mr-Apr '59. (MIRA 12:5)  
(PHOSPHORUS METABOLISM)

ILIN, V. S. (Prof.)

"Regulation hormonale de l'activite glucokinase du Foie."

report submitted for the 1st Intl. Symposium on Carbohydrate and Phosphorylated  
Glucides, 16-18 Sept 1960, Milan, Italy

Director, Inst. Exptl. Medicine, Acad. Med. Sci. USSR



IL'IN, V.S., prof.

Nikolai Vasil'evich Veselkin; on his eightieth birthday. Pat.fisiol.  
eksp.terap. 4 no.1:91-92 Ja-F '60. (MIRA 13:5)

1. Chlen-korrespondent ANS SSSR.  
(VESELKIN, NIKOLAI VASIL'EVICH, 1879--)

IL'IN, V.S.; SHANYGINA, K.I.

Hormonal regulation of the hexokinase reaction in the liver. Vop.  
med. khim, 6 no.3:291-300 My-Je '60. (MIRA 14:3)

1. Otdel biokhimii Instituta eksperimental'noy meditsiny Akademii  
meditsinskikh nauk SSSR, Leningrad.  
(LIVER) (HEXOKINASE) (DIABETES)  
(CORTISONE) (INSULIN)

IL'IN, V.S.; TRYUFANOV, V.F.

Incorporation of Methionine- $^{35}$  in liver proteins of rats with alloxan diabetes under conditions of a block or "circumvention" of the glucokinase reaction. Vop.med.khim. 6 no.4:386-389 J1-Ag '60.  
(MIRA 1419)

1. Department of Biochemistry, Institute of Experimental Medicine, the U.S.S.R. Academy of Medical Sciences, and Chair of Biochemistry of the S.M.Kirov Institute for Postgraduate Medical Training, Leningrad.

(DIABETES)

(METHIONINE)

(LIVER)

(PROTEIN METABOLISM)

ILIN, V. S., TRUFANOV, N. F., SHANGINA, K. I. (USSR)

"The Inclusion of <sup>35</sup>S Methionine into Proteins of the Liver  
in Block and -Circumvention- of the Glucokinase Reaction."

Report presented at the 5th International Biochemistry Congress,  
Moscow, 10-16 Aug 1961

LEVIN V. S.  
LEVIN, V.S.; TRYUFANOV, V.F.

Inclusion of methionine- $S^{35}$  into liver slices of rats with alloxan diabetes in a medium containing glucose or fructose. Vop. med. khim. 7 no. 1:82-85 Ja-F '61. (MIRA 14:4)

1. Chair of Biochemistry of the "S.M.Kirov" State Institute for Advanced Medical Training and Department of Biochemistry of the Institute of Experimental Medicine, Academy of Medical Sciences of the USSR, Leningrad.  
(DIABETES) (LIVER) (METHIONINE)

IL'IN, V.S.; NLYFAKH, S.A.

Mechanisms of cellular and central regulation of energy metabolism.  
Vest. AMN SSSR 16 no.11:47-55 '61, (MIRA 15:2)  
(CELL METABOLISM) (NERVOUS SYSTEM)

IL'IN, V.S.; TITOV, G.V.

Reproduction in vitro of the effect of insulin on the hexokinase activity of the liver. Dokl. AN SSSR 141 no.1:227-229 N '61.

(HHA 14:11)

1. Institut eksperimental'noy meditsiny Akademii meditsinskikh nauk SSSR. Predstavleno akademikom V.N.Chernigovskim.

(HEXOKINASE)

(INSULIN)

(LIVER)

IL'IN, V.S.; KILDEMA, L.A. [Kildeeva, L.]

Effect of cortisone and insulin on the activity of hexokinase  
in leucocytes. Vop. med. khim. 8 no.4:374-376 31-Aug '62.

(MIRA 17:11)

1. Otdel biokhimii Instituta eksperimental'noy meditsiny AN  
SSSR, Leningrad i sektor biokhimii Instituta eksperimental'noy  
i klinicheskoy meditsiny AN Estonskoy SSR, Tallin.



IL'IN, V.S.; TITOVA, G.V.

Effect of insulin in vitro on the activity of hepatic glucose-6-phosphate dehydrogenase and glucose-6-phosphatase. *Biochimia* 28 no.6:987-991 N-D'63 (MIRA 17:1)

1. Department of Biochemistry, Institute of Experimental medicine, Academy of Medical Sciences, Leningrad.

ZBARSKIY, Boris Il'ich [deceased]; IVANOV, Il'ya Il'ich;  
MARDASHEV, Sergey Rufovich; IL'IN, V.S., red.

[Biological chemistry] Biologicheskaya khimiya. 4. izd.,  
1 spr. i dop. Leningrad, Meditsina, 1965. 519 p.  
(MIRA 18:6)

GROMOVA, K.G.; IL'IN, V.S.

Identification of glucokinase in the adipose tissue of the epididymis of rat testicle and the stimulating effect of insulin on its synthesis. Biokhimiia 30 no.4:752-758 J1-Ag '65. (MIRA 18:8)

1. Otdel biokhimi Institute eksperimental'noy meditsiny AMN SSSR, Leningrad.

IL'IN, V. S., CAND AGR SCI, "INTERVARIETAL HYBRIDS OF  
CORN IN OMSKAYA OBLAST." OMSK, 1961. (AUTHOR'S ABSTRACTS  
OF DISSERTATIONS<sup>g</sup> PRESENTED AT OMSK AGR INST IM S. M. KIROV).  
(KL, 3-61, 225).

[L'IN, V.S.]

10 июня  
(с 10 до 18 часов)

Ю. Е. Пурман

Новый метод определения разности температур  
по разности тепловых потоков.

В. Н. Топкин

К вопросу о механизме диффузионной деформации.

В. Г. Шенк

Свойства полимеров с пространственной  
структурой.

10 июня  
(с 10 до 22 часов)

Г. Л. Фролов

Определение коэффициента диффузии в  
полимерах.

А. Н. Шенк

Новый метод определения температуры  
разложения полимеров.

10

А. С. Косов

В. А. Гусев

Влияние условий концентрации на процесс  
разложения полимеров.

С. Н. Шенк

Анализ данных о разложении полимеров.

В. А. Косов

А. Н. Шенк

Свойства полимеров с пространственной  
структурой.

11 июня  
(с 10 до 18 часов)

В. С. Шенк

Влияние температуры на процесс  
разложения полимеров.

В. С. Шенк

Влияние температуры на процесс  
разложения полимеров.

В. С. Шенк

Влияние температуры на процесс  
разложения полимеров.

11

papers submitted for the Scientific Meeting of the Scientific Societies of  
Radio Engineering and Electrical Communications in A. S. Paper (1959), Moscow,  
8-10 June, 1959

IL'IN, V.S.

Variational method of calculating symmetrical tetrapolar inhomogeneities in rectangular wave guides. *Izv.vys.sheeb.sov.; radiofiz.*  
3 no.4:683-693 '60. (MIRA 13:9)

1. Saratovskiy gosudarstvennyy universitet.  
(Wave guides)

KARMILOVA, L.V.; YENIKOLOPYAN, N.S.; HALBANDYAN, A.B.; IL'IN, V.T.  
(Moskva)

Kinetics and mechanism of methane oxidation. Part 5:  
Constant rate of methane oxidation. Zhur. fiz. Khim.  
35 no.7:1435-1442 J1 '61. (MIRA 14:7)

1. Akademiya nauk SSSR, Institut khimicheskoy fiziki.  
(Methane) (Oxidation)

L 10716-66 EWT(d)/EWT(1)/EPF(n)-2/EWP(1) LJP(c) BB/kw/X  
 ACC NR: AF5028515 SOURCE CODE: UR/0286/45/000/020/0098/0098  
 AUTHORS: <sup>44, 55</sup> Il'in, V. V.; <sup>44, 55</sup> Vasil'yev, G. I.; <sup>44, 55</sup> Novikov, I. B.; <sup>44, 55</sup> Tsvetkova, R. P. 70  
 ORG: none 3

TITLE: Analog-digital device for simulating problems of nonstationary heat conduction. Class 42, No. 175750

SOURCE: Byulleten' izobreteniy i tovarnykh znakov, no. 20, 1965, 98

TOPIC TAGS: <sup>21, 44, 55</sup> temperature simulation, analog digital converter, heat conduction 161, 44

ABSTRACT: This Author Certificate presents an analog-digital device for simulating nonstationary heat conduction. The device contains a network of ohmic resistances and a device for setting the boundary conditions. To increase the accuracy of simulation, the device contains a code-to-analog converter for each node, which operate alternately in a storage and a voltage setting mode. There is also one analog-to-code converter common to all the nodes, which is connected sequentially through an electronic switch to the nodes of the network.

Card 1/2

UDC: 681.14.001.572

L 10716-66  
 ACC NR: AF5028515

Codes proportional to the voltages at these points are supplied to the inputs of the code-to-analog converters operating in the storage mode.

SUB CODE: 09/

SUBM DATE: 18Jul64

Card 2/2



8/044/63/100/002/030/050  
NOV/1125

AUTHORS: Il'in, V.V., Borshchev, V.B., Rokhlin, P.Z.

TITLE: Can a machine think? (Survey of some viewpoints). (In order of discussion)

PERIODICAL: Referativnyy zhurnal, Matematika, no. 2, 1963, 44, abstract 27213 (Tr. Kazansk. aviats. in-ta, 1961, 1961, no. 55, 65 - 80)

TEXT: In the authors' opinion, among a number of philosophical problems of cybernetics under discussion one problem which was raised and given differing solutions by many cyberneticians requires further discussion. "The problem deals with the possibility or impossibility of carrying out an analogy between a computer and the brain to the point of admission (or negation) that there arises a property in a computer identical or close to consciousness or, more clearly, to thought. Can a machine think? - This clear question produced two mutually exclusive viewpoints: yes or no". The authors analyze the basic arguments in support of the impossibility in principle of designing a thinking machine: the algorithmic undeterminability of certain problems, the nonreducibility of think-

Card 1/2

Can a machine think? ....

S/044/03/00/002/040/050  
A050/A125

ing, as a special form of motion of matter, to the physical form of motion of matter (to a lower form); the impossibility of simulating the subjective psychological universe of man; in the authors' opinion none of these arguments can be admitted as conclusive. "On the basis of the absence of contradiction to the idea of the possibility of designing a thinking machine by any laws of science or philosophy, this idea is admissible as a fully fledged scientific hypothesis. Future experience will solve the question of the truth of this hypothesis". In conclusion, the authors develop the thought that "in a whole number of works in the last few years an attempt is present to comprehend the changes introduced in the content of attributes of matter by the science of the twentieth century. .. the so-called ontological nongeoctrism". Carrying out an analogy between quantum theory, the theory of relativity, the authors consider that cybernetics apparently is the science which gives us the first foundation for discovering the elements of geocentrism in our consciousness and knowledge", since, "by deciding in the affirmative the problem of the possibility of creating a thinking machine we thus admit that there can exist not just a unique highly organized material system (the brain) in which consciousness arose ... that other forms of highly organized material systems in which consciousness arises are possible and exist."

Card 2/2 [Abstracter's note: Complete translation]

A.M. Kondratov

FRIMER, A.I.; ZAYTSEV, P.V.; IL'IN, V.V.; NITEKHIN, Ye.P.

Apparatus for thermal and cathodic atomization and etching of  
metals in a gas discharge. Zav.lab.22 no.2:238-240 P. 1956.  
(Metallography--Apparatus and supplies) (MIRA 9:6)

80V/48-23-2-1/20

21(7)

AUTHORS:

Bashilov, A. A., Il'in, V. V.

TITLE:

The Coefficients of the Internal Conversion of Some Nuclear Transitions in  $As^{75}$  (Koeffitsiyenty vnutrenney konversii nekotorykh yadernykh perekhodov v  $As^{75}$ )

PERIODICAL:

Izvestiya Akademii nauk SSSR. Seriya fizicheskaya, 1959, Vol 23, Nr 2, pp 154-158 (USSR)

ABSTRACT:

In a previous paper (Ref 4) the authors determined the coefficients of the internal conversion (CIC) of some nuclear transitions in  $As^{75}$ , that is to say, they determined the absolute value

$$\alpha_K = g \tau \frac{n_e}{n_\gamma}$$

by measuring the number of conversion electrons  $n_e$  by means of the magnetic spectrometer and by measuring the number of photoelectrons  $n_\gamma$  struck out of a calibrated target by use of one and the same radioactive sources.  $\tau$ , the photoelectric absorption coefficient and  $g$ , the energy function are the parameters of the apparatus, which were determined with

Card 1/3

80V/48-23-2-1/20

The Coefficients of the Internal Conversion of Some Nuclear Transitions in  $As^{75}$

especially calibrated samples and  $\gamma$ -rays with well known CIC. The target was calibrated with  $\gamma$ -rays of  $ThB$ ,  $Au^{198}$  and  $Cs^{137}$  (Fig 1). The authors determined the CIC for nuclear transitions in  $As^{75}$  with the energies  $E_\gamma = 265, 280, 305$  and  $401$  kev, which correspond to direct transitions from these energy levels into the ground state of  $As^{75}$ . Due to the fact that the intensity of  $E_\gamma = 305$  and  $401$  kev is considerably weaker as compared to  $E_\gamma = 265, 280$  kev, two radiation sources of different thickness with  $\sigma = 0.1$  mg/cm<sup>2</sup> and  $\sigma = 0.5$  mg/cm<sup>2</sup> were used. Figures 2, 3, 4 and table 1 contain the experimental data for  $\tau$ ,  $g$ ,  $n_e/n_\gamma$  and  $\alpha_K$ . For comparative data of other authors see table 2. There is a deviation by about 10%. The determination of the transition types resulting from a comparison of the theoretical and experimental values of  $\alpha_K$  is given in table 3. For the energies  $E_\gamma = 401, 305, 280, 265$  kev spin and parity of the individual levels were determined to be:  $5/2^+, 9/2^+, 5/2^-, 3/2^-$  (ground state); corresponding transitions:

Card 2/3